

Appendix 'B'

Poor outcomes in Lancashire, as at August 2017

Each of the following outcomes is significantly worse than the England rate.

Overarching indicators

- Life expectancy at birth in males and females;
- Healthy life expectancy in males.

Wider determinants

- Density of fast food outlets;
- Gap in the employment rate between those with a learning disability and the overall employment rate;
- Gap in the employment rate between those with a long-term condition and the overall employment rate;
- Adults in contact with mental health services in employment;
- Adults in contact with mental health services in settled accommodation;
- Unpaid carers (CYP and adults);
- GCSE achieved 5A*-C including English and Maths with free school meal status;
- Hospital admissions for violent crime (including sexual violence).

Start well

- Breastfeeding initiation;
- Proportion of five year old children free from dental decay;
- Infant mortality;
- Admission for diabetes for CYP aged under 19 years;
- A&E attendances, 0-19 years;
- Population vaccination coverage - DTaP/IPV/Hib, PCV & booster, Hib/MenC booster, MMR, Flu;
- Newborn bloodspot screening coverage;
- Newborn Hearing screening coverage;
- Under 18 and 16 conceptions;

Appendix 'B'

- Hospital admissions caused by unintentional and deliberate injuries (0-14 years and 15-24 years);
- Persons admitted to hospital for alcohol-specific conditions (under 18's);
- Hospital admissions due to substance misuse (15 to 24 years);
- Hospital admissions for mental health conditions;
- Admissions of babies under 14 days;
- Hospital admissions self-harm (10 to 24 years);
- Children in care;
- Children killed and seriously injured (KSI) on England's roads;
- Looked after children aged under 18 years.

Live well

- Proportion of people meeting recommended 5 a day on a usual day (adults);
- Percentage of physically active/inactive adults;
- Excess weight in adults;
- Admission episodes for alcohol-related admissions;
- Deaths from drug misuse;
- Emergency hospital admissions for intentional self-harm;
- Self-reported wellbeing - low satisfaction score, low worthwhile, low happiness and high anxiety;
- Killed and seriously injured (KSI) casualties on England's roads;
- Breast cancer screening coverage;
- Abdominal Aortic Aneurysm screening coverage;
- Adults in contact with secondary mental health services;
- Cumulative percentage of the eligible population aged 40-74 years offered an NHS Health Check.

Age well

- Mortality rate from causes considered preventable;
- Under 75s mortality rate from cardiovascular disease;
- Under 75s mortality rate from cancer;
- Under 75s mortality rate from liver disease;
- Under 75s mortality from respiratory disease;

Appendix 'B'

- Mortality rate from communicable diseases including influenza;
- Suicide rate;
- Emergency readmissions within 30 days of discharge from hospital;
- Preventable sight loss;
- Health-related quality of life for older people;
- Permanent admissions to residential and nursing care homes, 65 years and above;
- Proportion of older people, 65 years and above still at home 91 days after discharge from hospital;
- Proportion of older people offered reablement services following discharge from hospital;
- Delayed transfers of care per month;
- Reducing avoidable emergency admissions;
- Fuel poverty;
- People aged 65 years and above receiving winter fuel payments;
- People supported through Adult Social Care Services who have had a review of their needs.